

BRACHYURA (CRUSTACEA, DECAPODA)

By D. J. G. GRIFFIN and J. C. YALDWYN*
Australian Museum, Sydney

Abstract

The Survey collected 1023 specimens of Brachyura belonging to 29 species and 10 families. Seven species were taken by the Portland Pier Survey in 1963, five of which are also represented in the Port Phillip Survey collection. Only four of the 38 species known from Western Port are represented in the collection. The majid *Paratymolus latipes* and the xanthid *Pilumnus acer* are recorded from Victoria for the first time; previous records of the graspid *Cyclograpsus audouinii* from Victoria are doubtful. Seventeen species known from Port Phillip are not represented in the collection. All are typically cool temperate species well known from SE. Australia. Four species of *Pilumnus* were represented in the collections and these are compared in detail with other SE. Australian *Pilumnus* species. Most abundant in Port Phillip are *Halicarcinus ovatus* and *H. rostratus* (Hymenosomatidae), *Notomithrax minor* (Majidae), *Ebalia (Phylyxia) intermedia* (Leucosiidae), *Litocheira bispinosa* (Goneplacidae), *Pilumnus tomentosus* and *P. monilifer* (Xanthidae), *Nectocarcinus integrifrons* and *Carcinus maenas* (Portunidae) and *Pinnotheres pisum* (Pinnotheridae). The majority of the species are found on the sandy areas around the edge of the Bay, particularly in the W. areas; no species was taken in the central deeper parts of the Bay. Ovigerous females of most species were collected in late summer. Parasitism by sacculinas was small and confined to two species of *Pilumnus*.

Introduction

The Survey was carried out over a period of six years (1957-63) and 317 stations were worked during this period (Macpherson and Lynch 1966). More than 1000 specimens of crabs were collected during the Survey and these form the subject of this report. A superficial collection of the fauna of Portland Harbour, on the Victorian coast near the S. Australian border, was made at the Occan Pier on 9-10 June 1963 (Jeanette E. Watson, pers. comm.). One pile below the pier was taken as an average and a swathe was cut from top to bottom (32 ft). The material was sealed immediately in plastic bags. Bryozoa colonies were broken up for enclosed fauna, ascidians were closely examined and crustaceans removed. The specimens from the Portland Survey and a small collection from Western Port are treated together with the specimens from the Port Phillip Survey.

In a checklist of the Brachyura of Victoria

*Present address: Dominion Museum, Wellington, New Zealand.

published near the beginning of this century, Fulton and Grant (1906b) listed 37 species from Port Phillip and 38 species from Western Port; 24 of these species were listed as occurring in both areas. The following species listed by them are at present known by specific names other than those used in their list (current name in brackets): *Gonatorhyuchus tunidus* (*Paramithrax barbicornis*), *Halinus truncatipes* (*Naxia spinosa*), *Leptomithrax australiensis* (*L. gaimardii*), *Pilumnus lanatus* (*P. etheridgei*), *Pilumnus pilosa* (*Heteropilumnus fimbriatus*), *Lioxantho haswelli* (*Megametope rotundifrons*), *Ovalipes trimaculatus* (*O. australiensis*), *Cyclograpsus punctatus* (*C. granulatus*). A number of other species is now placed in genera other than those used by Fulton and Grant. Sayce (1902) earlier gave a list of dredged Brachyura from Port Phillip; the species were identified by F. E. Grant. Several are not included in the list given by Fulton and Grant (1906b) and therefore we consider that Grant thought those earlier identifications to be in error.

Ward (1929), in a popular article on the

crabs of Port Phillip, dealt with 21 species. All but four were ones listed by Fulton and Grant: *Paramithrax minor* (currently placed in *Notomithrax*), *Nectocarcinus tuberculatus*, *Paragrapsus quadridentatus* (listed by Fulton and Grant as *Casmagnathus* (sic) *quadridentatus* from Bass Strait and Victorian coast) and *Petalomera lamellata* (*P. lateralis* in caption to figure) were the additional species. Several species were discussed under different names from those used by Fulton and Grant.

The present report deals with 31 species—29 from Port Phillip, eight species from the Portland area and four from Western Port. References are given for each species, one to the original description, one to the most recent treatment of the species and one or more to any other which discusses the species in detail or provides an adequate illustration. The total number of males and females (including ovigerous females), the size range (in mm) and the size of the smallest ovigerous female are given. The size given is the greatest width across the carapace (abbreviated as c.w.) or the greatest length of the carapace (c.l.) and is exclusive of spines except in the majids and hymenosomatids where the length of the rostrum is included and in the portunids where the lateral 'teeth' are included. Measurements were made to the nearest 0.1 mm with dial calipers. The localities from which specimens were taken by the Survey are listed by area number (with station number in brackets); each station number is followed by the number of specimens. Special attention is paid to those species which proved difficult to identify, where the Port Phillip collection permitted clarification of inter-specific differences, especially *Nectocarcinus* and *Pilumnus* species. The months during which ovigerous females were taken are listed for each species. Data on infestation by sacculinas are also included.

Family DROMIIDAE

Petalomera lateralis (Gray)

Dromia lateralis Gray, 1831: 40.

Petalomera lateralis; Rathbun, 1923: 153. Hale, 1927: 111-112, Figs. 108-109. Griffin, in press.

MATERIAL: 1 ♂, 1 ♀, c.w. 6.5, 11.3 mm. Survey areas 26 (301) 1 (number of specimens), 58 (293) 1.

REMARKS: The ridges on the ambulatory legs in this species are low and rounded, the carapace bears a close pubescence and the anterolateral teeth of the carapace are low and broad. These features are among those distinguishing *P. lateralis* from *P. lamellata* (Ortmann). See Griffin in press.

DISTRIBUTION: E., S., W. Australia from Low Isles, off Port Douglas, Qld., through N.S.W., Vict., Tasm., S. Aust. to Nickol Bay, a little N. of N.W. Cape, W. Aust. Intertidal to a depth recorded as '80-120 fm'. Unconfirmed records from Japan and the Philippine Islands.

Petalomera wilsoni (Fulton and Grant)

Cryptodromia wilsoni Fulton and Grant, 1902b: 61, Pl. 9 (*Dromia wilsoni* in caption).

Petalomera wilsoni; Rathbun, 1923: 154-156, Pl. 42, fig. 1. Hale, 1927: 113-114, Fig. 111. Dell, 1968: 14-17, Figs. 5-7, Pl. 2 (pleopods 1-2).

MATERIAL: 1 ♂, 1 ♀, c.w. 34.6, 41.2 mm. Survey area 58 (150-154) 1. Portland Pier Survey: from several small bryozoans under fisherman's pier, 9 June 63, 1 specimen.

REMARKS: These two specimens agree well with the previous description of this species. The pits and ridges formed by the tomentum on the carapace are characteristic.

DISTRIBUTION: SE. Australia from Port Stephens, N.S.W., through Vict. and Tasm. to a little W. of Kingston, SE. South Australia. Intertidal to 470 fm. N. and central New Zealand; central and S. Japan; Natal to Algoa Bay, S. Africa.

Dromidiopsis excavata (Stimpson)

Dromidia excavata Stimpson, 1858: 239; 1901: 172. *Dromidiopsis excavata*; Rathbun, 1923: 146-147, Pl. 38. Hale, 1927: 110, Fig. 106.

MATERIAL: 1 ♂, 4 ♀ (2 ovig.), c.w. 18.8-42.4 mm, smallest ovig. ♀ 41.8 mm. Survey areas 10 (13-15) 1, 31 (10) 1 in red sponge, 59 (23) 2, 68 (220) 1.

REMARKS: These specimens agree with the brief description given by Rathbun. The transverse fringe of long hairs just behind the front of the carapace is very distinctive.

The ovigerous females were taken in September.

DISTRIBUTION: SE. and S. Australia from Port Stephens, N.S.W., through Vict. and Tasm.

to Nuyts Archipelago, Great Australian Bight, S. Aust. Intertidal to a depth recorded as '70-100 fm'.

Family LEUCOSIIDAE

Ebalia (Phlyxia) intermedia Miers

Ebalia (Phlyxia) intermedia Miers, 1886: 308. Pl. 25, fig. 2-2c. Tyndale-Biscoe and George, 1962: 74, Fig. 4.4 (pleopod 1).

Phlyxia intermedia; Hale, 1927: 198-199, Fig. 199.

MATERIAL: 43 ♂♂, 48 ♀♀ (5 ovig.), c.l. 6.2-13.3 mm, smallest ovig ♀, 9.8 mm. Survey areas 3 (202) 25, 6 (66-67) 2, 7 (208) 10, 9 (178, 180) and 19 (179-181) 9, 11 (195) 11, 13 (92, 94) 4, 16 (283) 8, 18 (308) 3, 22 (119) 1, 27 (284) 1, 31 (10) 2, 42 (38) 1, (288) 3, (289) 1, 43 (303) 1, 55 (256) 3, 68 (155) 2, (157) 3. Additional Material: Area 42, Indented Head shore coll., J.H.M., 16 Jan. 64, 1 specimen.

REMARKS: The specimens vary in the shape of the intestinal lobe and length of the intestinal spine. The margin of the lobe is sometimes weakly convex with the lateral angles obtuse and sometimes the lateral angles are acute and the posterior margin straight or very weakly concave. The intestinal spine is usually rather short and blunt but is sometimes quite sharp.

The ovigerous females were taken during March.

DISTRIBUTION: S. and W. Australia from Western Port, Vict., through Tasm. and S. Australia to Cottesloe, just N. of Fremantle, W. Australia. Intertidal to 33 fm.

Philyra laevis Bell

Philyra laevis Bell, 1855: 300, Pl. 32, fig. 7. Hale, 1927: 194-195, Fig. 195. Tyndale-Biscoe and George, 1962: 75, Fig. 4.7 (pleopod 1).

MATERIAL: 10 ♂♂, 7 ♀♀, c.l. 6.5-22.0 mm. Survey areas 6 (118) 1, 27 (Point Wilson shore coll., 28 April 1962) 3, 42 (Indented Head shore coll., 1 Mar. 1959) 1, 49 (236) 1, 58 (89) 8, 59 (36) 1. Addition material: French Is., Western Port Bay, 20 Jan. 1963, 2 specimens.

REMARKS: The small specimens (up to about 10 mm c.l.) have the carapace sparsely granular, the lateral margins possess a few small tubercles or granules between the anterior two of the three usual subacute lobes and the merus

of the chelipeds is finely granular, particularly on the outer face where some granules are larger than others. In larger specimens (c.l. 15 mm) only the anterior part of the carapace is obviously granular.

DISTRIBUTION: S. Australia from Western Port, Vict., through Tasm. and S. Aust. to Albany area, SW. Western Australia. Intertidal to 6 fm.

Philyra undecimspinosa (Kinahan)

Bellidilia undecimspinosa Kinahan, 1856: 128, Pl. 3, fig. 2.

Ebalia (Phlyxia) undecimspinosa; Whitelegge, 1900: 162.

Philyra murrayensis Rathbun, 1923: 136-137, Pl. 34. Hale, 1927: 195-196, Fig. 196.

Philyra undecimspinosa; Griffin, in press.

MATERIAL: 5 ♂♂, 10 ♀♀ (5 ovig.), c.l. 9.1-30.7 mm, smallest ovig. ♀ 14.2 mm. Survey areas 3 (202) 6, 6 (63-67) 1, 7 (205) 3, 13 (82-83) 1, (94) 2, 43 (251) 2.

REMARKS: These specimens agree well with Kinahan's original description and with Rathbun's description of *Philyra murrayensis*. See Griffin, in press.

The ovigerous females were taken during March.

DISTRIBUTION: SE. and S. Australia from off Newcastle, N.S.W., through Vict. to S. Aust. waters (no detailed localities). Intertidal to 40 fm.

Family HYMENOSOMATIDAE

Halicarcinus ovatus Stimpson

Halicarcinus ovatus Stimpson, 1858: 109; 1907: 146. Stebbing, 1900: 523, Pl. 36A. Hale, 1927: 117, Fig. 113.

MATERIAL: 152 ♂♂, 138 ♀♀ (46 ovig.), c.l. 2.1-9.2 mm, smallest ovig. ♀ 4.4 mm. Survey areas 3 (202) 13, 5 (51) 3, (165-169) 6, 6 (66) 2, (118) 9, (136) 28, 7 (206) 14, (208) 1, 9 (84) 4, 9 (178, 180) and 19 (179-181) 7, 10 (13) 5, (Point Cook shore coll. 12 July 1960 2 specimens. 11 (190) 7, 13 (83) 1, (92) 8, 14 (4) 2, (95) 2, 16 (142) 7, 17 (173) 1, 18 (61) 3, 22 (119) 1, 27 (41) 8, (138-139) 2, (284) 3, 28 (140) 2, 30 (278) 2, 31 (10) 7, (275) 1, 39 (43) 1, 40 (101) 1, 42 Indented Head shore coll. 1 Mar. 59) 1, (109) 14, (281) 3, 50 (228) 5, (233) 3, (238) 1, 51 (250) 1, (271) 8,

55 (35) 7, (39) 10, 58 (80-81) 2, (88) 12, (150-154) 2, 59 (25) 3, (36) 4 visits, 14 specimens, (224) 2, 60 (235) 1, 61 (239) 2, (242) 1, 62 (96) 1, 64 (163) 6, 67 (216-217) 2, 68 (155) 12, 69 (221) 1. Portland Pier Survey: From Bryozoa 10-15 ft, J.E.W. 10 June 63, 2 specimens; in ascidian, just below low water mark 10 June 63, 1 specimen; in ascidian and sponge 30 ft, J.E.W. 10 June 63, 4 specimens; in fold in ascidian 20 ft, J.E.W. 9 June 63, 1 specimen; with encrusting ascidian (*Ascidia sydneyensis* Stimpson) 24 ft, J.E.W., 10 June 63, 1 specimen; in ascidian 9 ft, J.E.W., 9 June 63, 2 specimens; in ascidians, J.E.W., 9 June 63, 1 specimen; with ascidians (*Ascidia sydneyensis* Stimpson) 10 ft, J.E.W., 7 June 63, 1 specimen; with ascidians *Cystodites dellechiaiei* (Della Valle) 10 ft, J.E.W., 9 June 63, 2 specimens; with ascidians *Herdmania momus* (Savigny) 15-30 ft, J.E.W., 9 June 63, 3 specimens; with ascidians *Sycozoa cerebritiformis* Quoy and Gaimard 24 ft, J.E.W., 9 June 63, 2 specimens; from Bryozoa under fisherman's pier, J.E.W., 7 June 63, 3 specimens.

REMARKS: Of the wide variation shown by this species, that involving the rostrum, the pubescence of the carapace and legs, and the development of the proximal tooth on the dactyl of the cheliped in the male, is particularly obvious in the series examined here. The three rostral lobes are usually distinctly separated from each other throughout their length, subequal in length and more or less parallel. In most of the specimens from Portland Pier, however, the lateral rostral lobes are distinctly outwardly directed and are slightly more widely separated from the medial lobe than in most of the specimens from Port Phillip. In one specimen (ovig. ♀, e.l. 6.5 mm from Area 5) the medial lobe is a little longer than the laterals. In a large ♂ (e.l. 9.2 mm from Areas 9 and 19) the three lobes are adpressed throughout their length, the laterals curving inwards to meet the medial lobe from the base to the tip. The carapace is usually naked but several specimens have the dorsum of the carapace covered by hairs. The proximal, apically truncate, tooth on the inner edge of the dactyl in the male is present, although very small, in males from a

carapace length of about 4 mm and is prominent in large males (about 6 mm and above). In a few cases there is no marked proximal gape between the dactyl and fixed finger. In two males (c.l. 5.2 from Area 16, and e.l. 6.2 mm from Area 18) there is no trace of a proximal tooth and the fingers are adjacent throughout their length. Males as small as 2.5 mm possess well-developed pleopods.

Ovigerous females were taken in all months except July, November and December when no samples of this species were collected.

DISTRIBUTION: SE., S, and W. Australia from Sydney, N.S.W., through Viet., Tasm. and S. Aust. to Woodman's Point, just S. of Fremantle, W. Aust. Intertidal to 33 fm.

Halicarcinus rostratus (Haswell)

Hymenosoma rostratus Haswell, 1882: 550.

Halicarcinus rostratus; Kemp, 1917: 247. Hale, 1927: 117-118, Fig. 114.

MATERIAL: 17 ♂♂, 25 ♀♀ (17 ovig.), c.l. 2.1-9.2 mm, smallest ovig. ♀ 4.4 mm. Survey areas 11 (190) 5, 12 (196) 3, 14 (175) 1, 18 (308) 1, 19 (305) 1, 20 (124) 3, 36 (74, 76-77) 3, 39 (43) 1, 42 (109) 1, 43 (263) 1, 53 (253) 1, 55 (149) 4, 61 (241) 8, 62 (96) 2, (243-244) 2, 63 (21) 1, (162) 2, 68 (155) 2.

REMARKS: In males of this species less than about 6 mm e.l. the fingers of the chelae are adjacent throughout their length and lack the distinctive tooth pattern found in larger specimens.

Ovigerous females are present in samples taken during December, February, March, April, June and July.

DISTRIBUTION: S. Australia from Western Port, Viet., to Kangaroo Island and S. Aust. waters (no detailed localities). Intertidal to 11 fm.

Family MAJIDAE

Paratymolus latipes Haswell

Paratymolus latipes Haswell, 1880: 303, Pl. 16, figs. 3-5.

Paratymolus latipes var. *quadridentatus* Baker, 1906: 107, Pl. 1, fig. 2. Hale, 1927: 123, Fig. 119.

MATERIAL: 1 ♀, c.l. 6.4 mm. Portland Pier Survey: From small Bryozoa under fisherman's pier, 9 June 63, 1 specimen.

REMARKS: This specimen for the most part agrees with previous descriptions under the name *P. latipes* and *P. latipes* var. *quadridentatus*. The three anterolateral spines on each side have one tubercle midway between each at a slightly higher level. The merus of the cheliped bears one spine midway along the dorsal edge, and there are four similar spines along the ventrolateral edge. The palm of the chela is short and distally high and the dorsal edge bears two short, distally-directed spines.

DISTRIBUTION: E., S. and SW. Australia from Port Denison, Bowen, Qd., through N.S.W., Vict. and S. Aust. to Cockburn Sound, near Fremantle, W. Aust. Subtidal to 27 fm. Unconfirmed record from Ponape, Micronesia. This species has not been recorded previously from Victoria.

Naxia deflexifrons (Haswell)

Microhalimus deflexifrons Haswell; 1880: 435, Pl. 25, fig. 2.

Naxia (*Microhalimus*) *deflexifrons*; McCulloch, 1913: 330, Pl. 10, figs. 1-4.

MATERIAL: 4 ♂♂, 1 ♀, c.l. 6.3-15.0 mm. Survey areas 58 (88) 2, (150-154) 2, 59 (36) 1.

REMARKS: The five small specimens agree in all features with the descriptions and figures given by Haswell and McCulloch. The prominent lobe medial to the supraorbital eave is not obvious in small specimens, and the anterolateral spine of the basal antennal article is usually directed slightly laterally rather than forwards as shown by McCulloch.

DISTRIBUTION: SE. Australia from Port Jackson, N.S.W., through Bass Strait to Port Phillip, Vict. Intertidal to 37 fm.

Naxia aurita (Latreille)

Pisa aurita Latreille, 1825: 140.

Naxia aurita; Hale, 1927: 129, Fig. 127. Balss, 1935: 120 (synon.).

MATERIAL: 9 ♂♂, 9 ♀♀, c.l. 15.0-41.5 mm. Survey areas 26 (300) 2, 40 (Clifton Springs intertidal coll., 1 Mar. 59) 1, (101) 1; 42 (Indented Head intertidal coll. 1 Mar. 59) 3, (108) 1, 50 (233) 2, 59 (214) 1, (224) 2, 60 (235) 2, 67 (216) 1. Additional material: Area 42, Indented Head shore coll., 16 Jan. 64, 2 specimens.

REMARKS: This reasonably large series, comprising mostly small specimens, indicates that the best single features distinguishing small specimens of this species from small specimens of *N. aries* (Guérin) (see Hale 1927: 127, Fig. 128) is the straight, stout rostral spines. In small specimens of the present species the marginal spines of the carapace are relatively long as in *N. aries*.

DISTRIBUTION: S. and SW. Australia from D'Entrecasteaux Channel, Tasm., and Port Phillip, Vict., through S. Aust. to Abrolhos Islands, W. Aust. Intertidal to 8 fm.

Naxia tumida (Dana)

Halimus tumidus Dana, 1852: 115, Pl. 4, fig. 2a-g. *Naxia tumida*; Hale, 1927: 128, Fig. 126. Balss, 1935: 121 (synon.).

MATERIAL: 2 ♂♂, 4 ♀♀ (1 ovig.), c.l. 9.1-15.8 mm, ovig. ♀, 15.8 mm. Survey areas 58 (150-154) 2, (290) 1, 59 (79) 1, (87) 2.

REMARKS: The anterior part of the lateral margin of the basal antennal article bears one or a few short spines in all specimens, a feature characteristic of this relatively small species of *Naxia*.

The ovigerous female was collected in May.

DISTRIBUTION: E. and S. Aust. from Moreton Bay, Qd., through N.S.W. and Vict. to Kangaroo Is. and St. Vincent's Gulf, S. Aust. Intertidal to 7 fm.

Notomithrax minor (Filhol)

Paramithrax minor Filhol, 1885: 3; 1886: 356, Pl. 40, figs. 4-5, 7.

Notomithrax minor; Griffin, 1966: 53-57, Figs. 10, 21-3, 4.

MATERIAL: 70 ♂♂, 85 ♀♀ (42 ovig.), c.l. 5.2-37.0 mm, smallest ovig. ♀ 11.3 mm. Survey areas 3 (202) 5, 5 (51) 2, (166, 168) 3, 6 (66) 8, (118) 1, (137) 9, 7 (123) 1, (204) 7, (207) 1, (208) 16, 9 (178, 180) and 19 (179, 181) 2, 10 (103) 10, 11 (125) 1, (190) 13, 12 (196) 1, 13 (82-83) 5, (92, 94) 14, (209) 2, 16 (143) 2, 17 (170) 1, 18 (308) 1, 19 (305) 1, 21 (115) 1, 23 (2) 4, (68, 70) 4, (71) 1, 26 (301) 1, 27 (41) 1, 28 (141) 1, 29 (107) 1, (287) 1, 30 (278) 1, 35 (121) 2, 36 (77) 1, 37 (40) 3, 42 (Indented Head intertidal coll., 31 Mar. 59) 1, (108-109) 1, 50 (233) 1, 55 (35) 3, (Half Moon Bay intertidal coll., 14 Jan. 58) 1, (39) 3, (149)

1, (256) 1, 61 (239) 2, (242) 1, 62 (243-244) 1, 63 (20) 4, (159) 2, (245) 4, 68 (158) 1, (220) 1.

REMARKS: The very large series agrees in all features with material from New Zealand previously reported on by Griffin (1966) and from Australia recorded by Rathbun (1918) and by Haswell, Whitelegge, Fulton and Grant and Grant and McCulloch as *Paramithrax peronii* (for synonymy see Griffin 1966). There are low spines in the midline of the carapace, the protogastric regions are smooth and the marginal spines are alternately large and small. Small specimens (less than about 15 mm c.l.) have the anterolateral lobe of the basal antennal article spinulate or crenulate laterally and in many cases very small specimens have secondary spinules on the marginal spines.

Ovigerous females were taken in January, March, April, May, July, October, November and December.

DISTRIBUTION: E. and SE. Australia from Port Curtis, Gladstone, Qd., through N.S.W. to Port Phillip, Vict., and Tasm. Subtidal to a depth recorded as '22-60 fm'. N. and S. New Zealand to 70 fm.

Leptomithrax gaimardii (H. Milne Edwards)

Paramithrax gaimardii H. Milne Edwards, 1834: 325. *Leptomithrax australiensis*; Hale, 1927: 135-136, Fig. 135.

Leptomithrax gaimardii; Griffin, 1963: 133-137, Figs. 1-6, Pls. 6-7.

MATERIAL: 9 ♂♂, 7 ♀♀, c.l. 7.0-147.0 mm. Survey areas 18 (307-308) 1, 20 (309) 1, 27 (49) 1, 42 (108-109) 1, 43 (303) 1, 50 (228) 1, 51 (271) 1, 58 (150-154) 1, 59 (227) 1, 68 (220) 1, 69 (100) 1, (221) 1. Portland Pier Survey: on bottom and lower piles 10 June 63, 2 specimens; from bottom below piles 10 June 63, 1 specimen; J.E.W. (no date), 1 specimen.

REMARKS: Almost all the specimens have the carapace and chelipeds spinous rather than tuberculatc. In all important features they agree with specimens previously reported on by various authors (see references Griffin 1963). This species is easily recognized by the orange, transversely oval, naked area at the junction of the ischium and mcrus of the third maxilliped endopod.

The figures given by Griffin (1963), drawn from photographs of the holotype, do not accurately show the spinules around the orbit; the eave bears a number of spinules close to the margin, the intercalated spine bears several short, sharp spinules anteriorly and posteriorly near its base and the postorbital lobe bears a short stout spine on the anterior upper border about 0.3 of its length from the base.

DISTRIBUTION: SE. and S. Aust. from Shoalhaven Bight, off Nowra, N.S.W., through Vict., Tasm. and S. Aust. to Oyster Harbour, Albany, SW. Western Australia. Intertidal to a depth recorded as '250-450 fm'.

Family PORTUNIDAE

Carcinus maenas (Linnaeus)

Cancer maenas Linnaeus, 1758: 627.

Carcinus maenas; Stephenson and Campbell, 1960: 80-82, Figs. 1A, 2A; Pl. 1, fig. 1; Pl. 5A.

MATERIAL: 5 ♂♂, 16 ♀♀, c.w. 13.7-36.9 mm. Survey areas 5 (Altona intertidal coll.) 1 specimen, 9 (84) 14, 42 (38) 2, 58 (89) 4.

REMARKS: The 21 specimens have a granular carapace and the dactyl of the fifth leg (swimming paddle) is extremely narrow as is typical of this species.

DISTRIBUTION: SE. Aust. from Mallacoota Inlet, Vict. (near N.S.W. border) to Port Phillip, Vict. Intertidal. N. Atlantic; unconfirmed records from Red Sea and Hawaii.

Nectocarcinus integrifrons (Latreille)

Portunus integrifrons Latreille, 1825: 192.

Nectocarcinus integrifrons; A. Milne Edwards, 1861: 406-407, Pl. 38. Hale, 1927: 152-153, Fig. 153. Stephenson and Campbell, 1960: 83-84, Fig. 2B; Pl. 1, fig. 2; Pls. 5B, 6A.

MATERIAL: 18 ♂♂, 19 ♀♀ (2 ovig.), c.w. 5.3-73.6 mm, smaller ovig. ♀ 28.4 mm. Survey areas 3 (203) 1, 5 (51) 1, (168) 2, 7 (123) 2, 9 (178, 180) and 19 (179-181) 4, 10 (14) 2, 11 (190-192) 2, 14 (95) 1, 17 (173) 2, 18 (60-61) 3, 27 (41) 1, 39 (45) 1, 40 (101-102) 2, 42 (108-109) 4, 50 (228) 1, 51 (250) 1, (271) 1, 55 (39) 3, 59 (224) 1, 63 (21) 1. Additional material: Area 42 (Indented Head intertidal coll., 16 Jan. 64) 3 specimens. Portland Pier Survey: from bottom under pier 32 ft, 9 June 63, 1 specimen.

REMARKS: The following features, typical of this species, are present in the material from

Port Phillip. The frontal margin is smoothly but weakly convex, sometimes with a minute medial notch in very small specimens (c.w. less than 20 mm), or a shallow medial emargination in larger specimens; in one specimen (52 mm ♂ from Portland Pier) the medial frontal notch is prominent and narrow. In moderately large specimens the front is bordered in dorsal view by a single row of numerous small rounded tubercles with similar tubercles, fewer in number, behind them; a second, ventral, row of tubercles separated from the first by a fringe of hairs, is visible only in frontal view. Above the antenna, the frontal margin is very shallowly concave and the junction with the orbital margin is usually smoothly rounded. The anterolateral teeth of the carapace are sharp spines; in medium sized specimens these bear tubercles around their bases and along the lateral surfaces. The protogastric regions each possess a low elevation bearing small tubercles, and in front of these are more tubercles on a weak elevation. Usually the larger, posterior pair of elevations are weakly convex posteriorly and declivous anteriorly. In two specimens (males, 28.0, 31.2 mm, Area 10) the posterior elevations are uniformly convex, the anterior ones weakly declivous. The inner margin of the carpus of the cheliped bears several small tubercles in a row from the tip of the main spine to the articulation with the chela; in very small specimens there is usually no trace of spines or tubercles. In a few specimens between 20 mm and 30 mm c.w. these tubercles are sharp and enlarged. The surface of the carapace is pubescent in most of the specimens and beneath this the carapace is weakly tuberculate.

The ovigerous females were collected in March and October.

DISTRIBUTION: SE., S. and SW. Aust. from Port Stephens, N.S.W., through Viet., Tasm., S. Aust. to Cockburn Sound, a little S. of Fremantle, W. Aust. Intertidal to 8 fm.

***Ovalipes australiensis* Stephenson and Rees**

Ovalipes bipustulatus; Hale, 1927: 147-148, Fig. 148. (Not *Platyonichus bipustulatus* H. Milne Edwards, 1834.)

Ovalipes australiensis Stephenson and Rees, 1968: 227-232, Figs. 1-4; Pls. 35, 39, 41-42.

MATERIAL: 2 ♀ ♀, c.w. 24.3-57.1 mm. Survey areas 59 (36) 1, 63 (Safety Beach intertidal coll., 22 Sept 63) 1 specimen.

REMARKS: Both specimens have the carapace anteriorly granulate dorsally and a pair of orange (in alcohol) spots posteriorly; both features are characteristic of this species.

DISTRIBUTION: E., S. and SW. Aust. from Wide Bay, just S. of Fraser Is., Qd., through N.S.W., Vict., Tasm. and S. Aust. to Rottneest Is. and Cottesloe, just N. of Fremantle, W. Aust. (Presumably 'Shark Bay' in Stephenson and Rees 1968: 231 is a *lapsus* for Geographe Bay.) Subtidal to about 33 fm (60 m). Lord Howe Is., Tasman Sea.

Family XANTHIDAE

***Actaea peronii* (H. Milne Edwards)**

Xantho peronii H. Milne Edwards, 1834: 392.

Xantho spinosus Hess, 1865: 132, Pl. 6, fig. 3 (leg only).

Actaea peronii; Rathbun, 1923: 107, Pl. 21, figs. 4-5. Hale, 1927: 159, Fig. 159.

MATERIAL: 4 ♂ ♂, 6 ♀ ♀, c.w. 5.1-15.9 mm. Survey area 59 (Portsea Pier intertidal coll., 22 Mar. 1960) 1 specimen. Portland Pier Survey: from several small Bryozoa, 9 June 63, 7 specimens; in Bryozoa 15 ft, J.E.W., 9-10 June 63, 4 specimens.

REMARKS: All specimens possess the very long spines on the dorsal surface of the ambulatory carpi and propodi typical of this species. The tubercles on the lateral parts of the carapace are very large and rounded and the tubercles on the central areas are much lower and more discrete in small specimens with a tendency to become separated by transverse grooves anteriorly so that the carapace has the appearance of bearing numerous short transverse grooves centrally.

DISTRIBUTION: SE. and S. Aust. from Port Stephens, N.S.W., through Vict. and Tasm. to Spencer Gulf, S. Aust. Subtidal to a depth recorded as '70-80 fm'. Unconfirmed record from Samoa.

Two so-called subspecies of *A. peronii* have been recorded from Australia—*A. p. squamosa* Henderson from Torres Strait by Calman (1900) and *A. p. occidentalis* Odhner from SW. Australia (Odhner 1925: 58); the present




Fig. 1—Left first pleopods of males of *Pilumnus acer* (a-c), HOLOTYPE, c.w. 20.4 mm, 60-80 mi W. of Eucla, Great Australian Bight (AM E.3178) and *P. etheridgei* (d-f), HOLOTYPE, c.w. 15.6 mm, 10 mi N. of Circular Head, Tasm. (AM E.6490). a, f, whole pleopod, abdominal aspect; b, d, tip, abdominal aspect; c, e, tip, sternal aspect.

series from Victoria appears to belong to the typical form of *P. peronii*.

***Pilumnus acer* Rathbun (Fig. 1a-c)**

Pilumnus acer Rathbun, 1923: 124-125, Pl. 29. Hale, 1927: 165-166, Fig. 166.

MATERIAL: 13 ♂♂, 4 ♀♀ (1 ovig.), c.w. 5.1-21.0 mm, ovig. ♀ 21.0 mm. Survey areas 5 (58) 1, 27 (41) 3, 58 (89) 1, 59 (intertidal coll., 20 Mar. 60) 4, (36) 3, 61 (37) 1, 66 (292) 1. Portland Pier Survey: from small Bryozoa under fisherman's pier, J.E.W., 9 June 63, 1 specimen; in Bryozoa 15 ft, J.E.W., 10 June 63, 1 specimen; from Bryozoa 15 ft, J.E.W., 9 June 63, 1 specimen.

REMARKS: The very long, simple hairs forming a sparse but obvious fringe just behind the front, together with the long, simple, curved anterolateral spines, smooth carapace and the presence of sharp spines on the ambulatory meri and carpi, but not propodi, distinguish this species from its southern temperate congeners (see Table 1). Very small specimens (up to about 8 mm c.w.) have tufts of long thick hairs on the dorsal surface of the carapace.

The ovigerous female was taken in December. The preserved eggs are large (1.3 to 1.5 mm in diameter) suggesting that abbreviated or direct development may take place in this species as has been recorded for the Australian *Pilumnus vestitus* and the two New Zealand species of *Pilumnus*, *P. novaezealandiae* and *P. lumpinus* (see Wear 1967).

DISTRIBUTION: S. Australia from Port Phillip, Vict., through S. Aust. to a little W. of Eucla, Greath Australian Bight, W. Aust. Subtidal to a depth recorded as '80-120 fm'. This species has not been previously recorded from Victoria.

***Pilumnus etheridgei* Rathbun (Fig 1d-f)**

Pilumnus lanatus; Fulton and Grant, 1906b: 18. (Not *Pilumnus lanatus* Latreille, 1825.)

Pilumnus etheridgei Rathbun, 1923: 117-119, Pl. 26. Balss, 1933: 27.

MATERIAL: 3 ♂♂, 3 ♀♀, c.w. 5.6-13.3 mm. Survey areas 58 (88) 3, (154) 1, 59 (36) 1, 66 (292) 1.

REMARKS: The characteristic features of this species are the sparse, moderately long, simple hairs which generally occur on the anterior half

to two-thirds of the carapace dorsally and on the chelipeds and ambulatories (the ventral half or slightly more of the outer surface of the palm of the major chela is naked or bears some very slender hairs), the presence of sharp spines on the ambulatory carpi and propodi dorsally and the apically very strongly recurved first pleopod in the male. Accessory spinules are seldom present on the anterolateral spines.

Even small male specimens (about 8 mm c.w.) can be distinguished from the similar *P. acer* by the shape of the first pleopod, which is apically rather weakly curved in the latter species, and by the absence of a fringe of long hairs close to the front of the carapace.

DISTRIBUTION: S. and SW. Aust. from Oyster Bay, E. Tasm., through Vict. and presumably S. Aust. (but no records available) to off Fremantle, W. Aust. Subtidal to 26 fm.

***Pilumnus monilifer* Haswell (Fig. 2d-f)**

Pilumnus monilifera (sic) Haswell, 1881: 543-544; 1882: 65, Pl. 1, fig. 3.

Pilumnus monilifer; Hale, 1927: 163, Fig. 163.

MATERIAL: 37 ♂♂, 38 ♀♀ (3 ovig.), c.w. 4.1-21.7 mm, smallest ovigerous ♀ 9.1 mm. Survey areas 5 (51) 3, (Point Cook intertidal coll. 12 July 60) 6 specimens; 6 (137) 4, 10 (14) 1, 11 (190) 1, 14 (95) 1, 26 (301) 4, 27 (41) 10, (138) 1, 28 (141) 3, 30 (278) 1, (279) 2, 31 (131-134) 3, 37 (40) 1, 39 (43) 1, 42 (281) 1, 55 (149) 1, 58 (80) 3, 59 (36) 3, 61 (37) 1, (239) 2, 62 (96) 1, 68 (155) 1, 69 (221) 1, (222) 1. Portland Pier Survey: in sponges 10 ft, J.E.W., 10 June 63, 1 specimen; from Bryozoa 10-15 ft, J.E.W., 10 June 63, 2 specimens; from Bryozoa 15 ft, J.E.W., 9 June 63, 2 specimens; from small Bryozoa under fisherman's pier, J.E.W., 4 June 63, 13 specimens.

REMARKS: Large specimens (greater than about 12 mm c.w.) of the present series agree with the type material of Haswell's species (in Australian Museum—see Griffin, in press) particularly in the following features: there are tubercles on the anterolateral subspiniform lobes; an elevated group of about six tubercles is situated close to the anterolateral margin; the chelipeds and legs are mostly covered by a very dense mass of very short, 'clubbed' hairs




Fig. 2—Left first pleopods of males of *Pilumnus rufopunctatus* (a-c), c.w. 20.6 mm, Cabbage Tree Bay, N.S.W. (AM P.860) and *P. monilifer* (d-f), c.w. 21.0 mm, rocky shore below low tide mark, Beaumaris, Port Phillip, Vict. (AM P.9231). a, f, whole pleopod, abdominal aspect; b, d, tip, abdominal aspect; c, e, tip, sternal aspect.

with expanded setose tops; the outer surface of the palm of the major chela is covered by tubercles for slightly more than the dorsal half, and the ambulatory carpi and propodi possess several tubercles more or less in a double row dorsally. The tubercles on the carapace, chelipeds, and legs in most cases have retained a red colour and barely show through the mat of short hairs. The tubercles on the outer surface of the palm of the major chela tend to cover more of the surface in females than in males. In addition, the first pleopod of the male is apically strongly recurved with short hairs along the medial and lateral surfaces in the distal third to one half and a group of very long hairs on the lateral surface of the apical curve. In most specimens the carapace bears several tufts of very stout, long hairs.

In the key to Australian *Pilumnus* species given by Rathbun (1923: 108-110), *P. monilifer* is placed in that group of species having the carapace covered only by short hairs mixed with long hairs, and is distinguished from others in that group (particularly *P. rufopunctatus* Stimpson) in having the anterolateral lobes 'capped by a cluster of granules'. Hale (1927: 162-163), in a key to S. Australian species based on Rathbun's key, states that *P. monilifer* differs from *P. rufopunctatus* in lacking bead-like granules showing through the pubescence. Thus, in both Rathbun's and Hale's keys, large specimens from the present series come out only with difficulty.

Initially, many of the small specimens (less than about 12 mm c.w.) in the present series (now assigned to *P. monilifer*) were separated as a distinct species on the basis of the absence of obvious tubercles from the carapace and ambulatories, the simple spiniform nature of the anterolateral lobes and the less strongly apically curved first pleopod in the males. In those specimens from areas where silt and clay make up the dominant fraction of the substrate, the carapace, chelipeds and ambulatories possess many tufts of long thick hairs. However, these small specimens were identified as *P. monilifer* when it was found, from examination of large series of other species of *Pilumnus*, that in general smaller specimens in this genus tend to be less tuberculate and spinous than adults

and that in small males the first pleopod is less strongly curved apically. It must be stressed that some specimens of *P. monilifer* as small as 9 mm c.w. have supplementary tubercles on the slopes of the anterolateral lobes and that most specimens above this size possess at least three tubercles in a group near the anterolateral border.

Two other species of south-eastern Australian *Pilumnus* possess the close pubescence of the 'clubbed' type found on *P. monilifer*—*P. rufopunctatus* and *P. fissifrons* Stimpson (see Table 1). These also agree in the general arrangement of hairs and tubercles on the outer surface of the palm of the major chela and the dorsal surfaces of the ambulatories, but differ in the tuberculation of the carapace in adults. However, *P. rufopunctatus* never possesses tufts of long thick hairs and in *P. fissifrons* the first pleopod in males lacks long hairs laterally near the apical curve. It is by these features that *P. monilifer* can be distinguished from its congeners, not really by the characters selected by Rathbun and incorrectly modified by Hale.

The ovigerous females were taken in June and September.

One specimen, a female, c.w. 8.8 mm, is infested with a *Sacculina*.

DISTRIBUTION: Southern Australia from Victorian, Tasmanian and S. Australian waters (no detailed range limits available). Intertidal to 10 fm.

Pilumnus tomentosus Latreille (Fig. 3a-c)

Pilumnus tomentosus Latreille, 1825: 125. Rathbun, 1923: 119-122, Pl. 27, figs. 1-2. Hale, 1927: 166. Fig. 167. Balss, 1933: 23 (part: not Pl. 3, figs. 14-15).

MATERIAL: 16 ♂♂, 34 ♀♀, c.w. 5.6-39.6 mm. Survey areas 5 (51) 1, (166) 1, 6 (66) 1, (137) 1, 7 (123) 2, 10 (13) 4, 11 (190) 3, 13 (94) 4, 14 (4) 2, 16 (142) 1, 17 (172) 1, 18 (308) 1, 26 (301) 3, 28 (286) 2, 35 (121) 3, 36 (77) 1, 39 (43) 1, 40 (101) 1, 47 (29) 1, 51 (250) 2, 62 (96) 3, 63 (20) 2, 64 (164) 3, 69 (100) 1, (222) 1. Additional Material: Areas 63 (1 mile 1.6 km, off Dromana on sandy bottom, dredged 3 fm 22 Sept. 62), 2 specimens; 68 (1.5 miles, 2.4 km, off Rye Pier 7 fm, T. Crawford 16 Mar. 63), 2 specimens.




Fig. 3.—Left first pleopods of males of *Pilumnus tomentosus* (a-c), c.w. 31.7 mm, 80-120 fm, 60-80 mi W. from Eucla, Great Australian Bight (AM P.3562) and *P. australis* (d-f), c.w. 10.6 mm, from intestine of Nannygai (fish) caught at Shellharbour, N.S.W. (AM P.8442). a, f, whole pleopod, abdominal aspect; b, d, tip, abdominal aspect; c, e, tip, sternal aspect.

REMARKS: This species is characterized mainly by the presence of long, simple hairs rather densely covering the carapace, chelipeds and legs, the presence of tubercles over all of the outer surface of the palm of the major chela—these tubercles tend to be pointed and longer, but sparser, dorsally—and by the lack of spines or tubercles on the carpi and propodi of the ambulatory legs. Larger specimens (above about 25 mm c.w.) usually have one to three spinules or tubercles on the posterior slopes of the anterolateral spines and about three spines on the dorsal surface of the carapace near the anterolateral border. Small specimens (less than 15 mm c.w.) often possess tufts or long thick hairs on the dorsal surface of the carapace and along the dorsal and ventral edges of the ambulatory legs and usually lack spines on the dorsal surface of the carapace and lack spinules on the anterolateral lobes.

Nine specimens ranging in size from 8.5-12.5 mm c.w., are infested with sacculinas.

DISTRIBUTION: SE. and S. Australia from off Newcastle, N.S.W., through Vict., Tasm. and S. Aust. to Albany in SW. Western Australia. From 3 fm to a depth recorded as '200-300 fm'. Extra-Australian records of *P. tomentosus* are now considered as referring to other species of this genus (see Griffin, 1970).

General remarks on SE. Australian *Pilumnus* species

Nine species of *Pilumnus* occur in SE. Australia. (Balss 1933: 11-13, has transferred the deep water *P. spongiosus* Nobili to *Planopilumnus*; *P. terraereginae* Haswell is described and figured elsewhere (Griffin, 1970)). Table 1 compares seven of these with respect to a number of characters. Four have been collected in the Port Phillip Survey and two others have




Fig. 4—Left first pleopod of male of *Pilumnus fissifrons*, c.w. 13.7 mm, among growth on hull of H.M.A.S. *Penguin* in dock, Cockatoo Island, Port Jackson, N.S.W. (AM P.8563). a, tip, abdominal aspect; b, tip, sternal aspect; c, whole pleopod, abdominal aspect.

been discussed in relation to *P. monilifer*. The seven species can be divided into two groups which are fairly easy to separate from each other: *P. monilifer*, *P. rufopunctatus* Stimpson (see Rathbun 1923: Pl. 24, figs. 3-4, and Takeda and Miyake 1968: 12-15, Fig. 3a-c, Pl. 1D) and *P. fissifrons* Stimpson (see Hale 1927: Fig. 164, Takeda and Miyake 1968: 15-17, Fig. 3d-f, Pl. 1A), share features of hairiness and ornamentation of the carapace, chelipeds and ambulatory legs and differ in these features from *P. tomentosus*, *P. acer*, *P. australis* Whitelegge (see Whitelegge 1900: Pl. 35, figs. 1-4) and *P. etheridgei*, which, however, differ from each other in ornamentation of the ambulatories. Within these groups separation can be difficult, especially with small specimens.

The only other south-eastern Australian *Pilumnus* is *P. senilanatus* Miers (see Rathbun 1923: Pl. 24, figs. 1-2, Takeda and Miyake 1968: 7-9, Fig. 1a-c, Pl. 1B) which occurs in S. Queensland and S. Australia. This differs from the other six in almost all of the features dealt with in the table: the first anterolateral lobe is rounded and the last two are blunt, all these lobes are covered by numerous small tubercles, the hairs on the carapace are in rows anteriorly, the tubercles on the carapace occur mainly near the front in the midline, where a prominent group covers a pair of postfrontal elevations, the subhepatic region is smooth, the hairs on the outer surface of the palm of the major chela are confined to a small proximal area and do not occur over the dorsal surface, on the minor chela the hairs form a dense mass on the outer surface but both the dorsal and ventral surfaces are smooth and the hairs on the ambulatories form a dense mass on the posterior surface of the propodi as well as along dorsal and ventral edges of all segments.

***Heteropilumnus fimbriatus* (H. Milne Edwards)**

Pilumnus fimbriatus H. Milne Edwards, 1834: 416.
Heteropilumnus fimbriatus; De Man, 1895: 533, 536.
 Hale, 1927: 168, Fig. 170. (Not *Pilumnus fimbriatus* Haswell, 1882 (= *Cryptocoeloma haswelli* Rathbun, 1923; see Balss 1933: 43).)
Pilumnus pilosus Fulton and Grant, 1906a: 7, Pl. 4, figs. 1-4.

MATERIAL: 5 ♂♂, c.w. 5.5-14.6 mm. Survey areas 7 (123) 1, 14 (95) 1, 22 (119) 1, 25 (299) 1, 42 (281) 1.

REMARKS: All specimens possess the very dense fringe of long silky hairs around the front and anterolateral borders, the four low lobes on the anterolateral margin and the smooth carapace typical of this species. In all specimens the outer faces of both chelae are covered by close-set small tubercles. On the minor chela these tubercles usually cover almost the whole of the outer surface of the palm leaving bare only a small area near the articulation of the dactyl. The major chela in most specimens is also densely tuberculate, the tubercles covering at least the proximal half of the outer surface of the palm. In one specimen (♂, c.w. 12.1 mm from Area 42), however, the tubercles on the major chela are restricted to a small proximal area near the articulation with the carpus, the remainder of the outer surface being smooth and bare; the minor chela in this specimen possesses tubercles ventrally and dorsally and the proximal tubercles extend on to the outer face to leave a larger area smooth and naked than in other specimens. Fulton and Grant's type material of *Pilumnus pilosus* (one 'co-type' in Australian Museum, a dry ♂, c.w. 11.3 mm, mounted on glass with printed label 'Cryptocoeloma pilosus Fult. & Grant' and registered as G.5907) resembles the Port Phillip Survey material fairly closely in the granulation of the chelae although the original description does not make this clear.

DISTRIBUTION: S. Australia from Western Port, Vict., through Tasm. to St. Vincent Gulf, S. Aust. Intertidal to 11 fm. Records under this name from tropical Australia and other Indo-pacific localities presumably refer to *Cryptocoeloma haswelli* Rathbun.

***Pilumnopeus serratifrons* (Kinahan)**

Ozius (?) *serratifrons* Kinahan, 1856: 118, Pl. 4, fig. 1.
Heteropanope australiensis Stimpson, 1858: 87; 1907: 64, Pl. 7, fig. 7.
Pilumnopeus crassimanus A. Milne Edwards, 1867: 228.
Pilumnopeus serratifrons; Haswell, 1882: 70, Pl. 2, figs. 1a-b. Balss, 1933: 34 (synon.). Dell, 1968: 19-20, Figs. 1-4 (pleopod 1), Pl. 3.
Heteropanope serratifrons; Hale, 1927: 161, Fig. 162.

MATERIAL: 5 ♂♂, 1 ♀ (ovig.), c.w. 16.9-26.9 mm, ovig. ♀ 16.9 mm. Survey areas 27 (Point Wilson intertidal coll. 28 Apr. 62) 1 specimen, 42 (38) 2. Additional material:

Table 7. Comparison of south-eastern Australian species of *P. limbus*

Structure	Character	<i>P. maculifer</i>	<i>P. rufocinctatus</i>	<i>P. flavipes</i>	<i>P. tomentosus</i>	<i>P. cyan</i>	<i>P. emarginatus</i>	<i>P. quadratus</i>
Anterior-lateral lobes	form	short, subcapitiform external orbital angle blunt	short, stout, curved spines	short, stout, spiniform	moderately long, stout spines external orbital angle short	long spines, external orbital angle short	moderately long, stout spines	very long, stout spines, no spine on external orbital angle
	ornamentation	1-3 tubercles on medial or posterior slopes	1 or 2 tubercles on posterior slopes	none	1-3 spinules on posterior slopes	none	1-2 spinules sometimes in front or behind first two	none
Coropoda	hairs	dense mat of short 'clubbed' hairs, especially anteriorly, mixed with tufts of long, stout hairs	dense mat of short 'clubbed' hairs mixed with long hairs	dense mat of 'clubbed' hairs and several tufts of long stout hairs	moderately long, simple, stout longer and more numerous anteriorly	very long, simple, covering whole of surface, forming a fringe just behind front	moderately long, simple, not especially dense, sparse posteriorly	simple hairs of various lengths, sparse posteriorly, longer and more dense anteriorly, some short 'clubbed' hairs posteriorly
	ornamentation	one main group of 2-10 tubercles in curved row near anterolateral border, other tubercles near last anterolateral lobe and anteriorly submedially	several groups of a few tubercles near anterolateral border, near to last anterolateral lobe and on gastric regions medially	a few tubercles or spinules near anterolateral border and submedially anteriorly	a group of 1-3 spines near anterolateral border and sometimes one or more behind and submedially	none, coropoda smooth	none, coropoda smooth	none, coropoda smooth
Frons	ornamentation	closely-spaced, small tubercles	closely-spaced, small tubercles	closely-spaced tubercles	small tubercles	sometimes with small tubercles	close-spaced tubercles	about 4 slender spinules on each side, lateral one widely separated from others
Orbita	ornamentation	closely-spaced, blunt tubercles	closely-spaced, pointed tubercles	a few tubercles dorsally near external angle, closely-spaced tubercles ventrally	blunt or pointed tubercles dorsally and ventrally	smooth dorsally, spinules ventrally	stout spinules dorsally and ventrally	unpaired dorsally, ventrally with minute, widely spaced tubercles and one long spine medially
Subcapitulum region	ornamentation	a few small tubercles, one or more enlarged	1 or 2 tubercles	a few small spinules	several tubercles and 1-4 spines	several tubercles and spinules	several tubercles and spinules	minute spinules and tubercles
Major chela palm - outer surface	hairs	dense mat of short 'clubbed' hairs, ventral $\frac{1}{2}$ or more naked	dense mat of short 'clubbed' hairs mixed with some longer simple hairs dorsally, ventral $\frac{1}{2}$ or more naked	scattered long, simple hairs and dense mat of short 'clubbed' hairs, ventral $\frac{1}{2}$ or more naked	moderately long, simple, stout, more dense dorsally, ventral $\frac{1}{2}$ or less naked	very long, simple, ventral $\frac{1}{2}$ or more naked	moderately long, simple, not especially dense, leaving just more than ventral $\frac{1}{2}$ naked	very long, simple hairs, ventral $\frac{1}{2}$ or more naked
	ornamentation	closely-spaced, round, blunt tubercles, ventral $\frac{1}{2}$ or more smooth	closely-spaced, round, blunt tubercles, pointed dorsally, ventral $\frac{1}{2}$ or less smooth	closely-spaced round, blunt tubercles, ventral $\frac{1}{2}$ or more smooth	stout tubercles covering whole surface, blunt ventrally, sharper, longer and less dense dorsally	large, pointed tubercles, dorsal ones sometimes longer and curved, ventral $\frac{1}{2}$ or more smooth	small, closely spaced, stout spines covering almost all of surface, longer and sharper dorsally	closely spaced, stout spines on dorsal $\frac{1}{2}$ - $\frac{2}{3}$, small and blunt towards ventral edge, longer, sharper dorsally
Minor chela palm - outer surface	hairs	dense mat of short 'clubbed' hairs, ventral $\frac{1}{2}$ or more naked	dense mat of short 'clubbed' hairs mixed with some short simple hairs, ventral $\frac{1}{3}$ or less naked	scattered long, simple hairs and dense mat of short 'clubbed' hairs, ventral $\frac{1}{2}$ or more naked	moderately long, simple, covering whole of surface, more dense dorsally	some simple, very long dorsally, shorter ventrally, covering whole of surface	moderately long, simple, not especially dense, leaving ventral $\frac{1}{2}$ - $\frac{2}{3}$ naked	very long, simple hairs covering all of surface, generally sparse but more dense dorsally
	ornamentation	closely-spaced, round blunt tubercles, ventral $\frac{1}{2}$ or more smooth	tubercles covering whole of outer surface, pointed dorsally	small blunt tubercles, ventral $\frac{1}{2}$ or less smooth	stout, pointed tubercles more or less in rows, over whole surface, less dense dorsally	tubercles ventrally and spines dorsally more or less in rows, longer and curved dorsally	small, closely spaced, stout spines covering almost all of surface, longer and sharper dorsally	stout spines covering all of surface, longest dorsally
Ambulatory legs	hairs	dense mat of short 'clubbed' hairs dorsally, longer simple hairs dorsally and ventrally	dense mat of short 'clubbed' hairs dorsally, longer simple hairs dorsally and ventrally	dense mat of short 'clubbed' hairs on dorsal and posterior surfaces, long, stout and slender hairs dorsally and ventrally	simple, long, dense dorsally, shorter on posterior and anterior surfaces	very long, simple, straight, longer and more dense dorsally	long, simple, rather sparse	simple hairs on all surfaces of all segments, dorsally longer and more dense
Ambulatory manus	ornamentation	dorsal edge smooth or with a few very low tubercles	dorsal edge with a few curved spinules	smooth or with a few tubercles	sometimes a few spinules dorsally	up to 5 short, curved spines on dorsal edge	a few tubercles dorsally	4 or more spines dorsally, longer distally, some short spinules ventrally
Ambulatory carpus	ornamentation	several blunt tubercles more or less in a double row	several blunt to spiniform tubercles dorsally more or less in double row	small, blunt tubercles dorsally more or less in double row	lacking spines or tubercles	about 3 long, straight or weakly curved spines dorsally	up to 6 long spines dorsally	3 long spines dorsally
Ambulatory propodus	ornamentation	several blunt tubercles more or less in a double row	several blunt to spiniform tubercles dorsally more or less in double row	small, blunt tubercles dorsally more or less in double row	lacking spines or tubercles	lacking spines or tubercles	about 6 long spines more or less in double row	none